

Comparing Ultraviolet and H α Star Formation Rates

Eric F. Bell and Robert C. Kennicutt, Jr.

Steward Observatory, 933 N. Cherry Ave., Tucson, AZ 85721, U.S.A.

Abstract. We have used a sample of 43 star-forming galaxies imaged in the far-ultraviolet (FUV) by the UIT supplemented with 33 galaxies observed by FAUST, to explore the consistency of UV and H α derived star formation rates (SFRs). We find, even before correction for dust, that UV and H α SFRs are quantitatively consistent for low-luminosity galaxies, and that higher luminosity galaxies have H α SFRs a factor of 1.5 higher than their UV SFRs: this reflects the influence of dust. Our results are consistent with a scenario where the UV dust extinction is a factor of $\lesssim 2$ larger than the H α dust extinction in a given galaxy, and that there is over a 4 magnitude range of H α dust extinctions which correlate loosely with galaxy luminosity such that low luminosity galaxies tend to have lower extinctions than their higher luminosity counterparts.

1. The Data

We have selected a sample of 43 galaxies with FUV detections or upper limits and H α luminosities from the sample of spiral galaxies imaged using the Ultraviolet Imaging Telescope (UIT). Magnitudes at an effective wavelength of 1567 Å were derived; from external comparisons, we estimate that our magnitudes are accurate to at worst 0.4 mag, and typically 0.2 mag. Magnitudes for a further 33 galaxies observed by FAUST (Deharveng et al. 1994) at 1650 Å were also included in the analysis as a check.

H α luminosities are primarily taken from the literature, with the addition of some unpublished CCD data taken by the authors. When appropriate, the total flux was corrected for [N II] following Kennicutt (1983). To determine the importance of dust, we also examined far-infrared (FIR) and thermal radio continuum luminosities. The FUV to FIR ratio is a rough probe of dust attenuation. The H α to thermal radio ratio is a direct probe of the attenuation of the H α light as the two radiations are produced by the same thermal electrons: the only disadvantage is that the thermal radio luminosity is difficult to measure accurately in most cases.

2. Results

In panels a and b of Fig. 1, we can see that the H α SFRs are a factor of ~ 1.5 larger than the FUV SFRs for high luminosity galaxies, and comparable for lower luminosity galaxies. This is consistent with a scenario in which both the




Figure 1. **a)** Comparison of the UV and H α SFRs (calibrated using Kennicutt 1998) for UIT ($\bullet$) and FAUST ($\circ$) galaxies. The solid line denotes equal SFRs, and the dotted line denotes the least squares fit to the data. **b)** H α to UV SFR ratio with V band absolute magnitude. Symbols as in a): the dashed line denotes the trend in H α to FUV SFR ratio expected because of metallicity effects. **c)** H α attenuation determined from the radio against the FIR to FUV luminosity ratio. The solid line denotes the expected correlation if the H α attenuation is half of the FUV attenuation: the dotted lines show the correlation if the H α attenuation is 1/4 of (lower line) or equal to (upper line) the FUV attenuation.

H α and FUV radiation suffer attenuation (where the FUV attenuation is larger than the H α attenuation), and that the overall amount of attenuation varies as a function of galaxy luminosity (e.g. Tully et al. 1998).

We explore the attenuation more directly in panel c, where we plot the H α attenuation determined from the H α to thermal radio ratio against the FIR to FUV luminosity ratio (a tracer of FUV attenuation, modulo uncertainties from older stellar population heating of the dust). The H α attenuations are clearly correlated with the FUV attenuations, indicating a real spread in overall spiral galaxy attenuation. Furthermore, the data are consistent with a H α attenuation $\sim 1-2$ times lower than the FUV attenuation (simple models predict a factor of 4 offset: models that take into account the different distribution of FUV and H α emitting sources predict a factor of 2 offset). A more thorough discussion of these results is presented by Bell & Kennicutt (2000).

Acknowledgments. This work was supported by NASA LTSA grant NAG5-8426 and NSF grant AST-9900789.

References

- Bell, E. F. & Kennicutt, R. C., Jr. 2000, submitted to ApJ
- Deharveng, J. M., et al. 1994, A&A, 289, 715
- Kennicutt, R. C., Jr. 1983, ApJ, 272, 54
- Kennicutt, R. C., Jr. 1998, ARA&A, 36, 189
- Tully R. B., et al. 1998, AJ, 115, 2264